

## Accepted Manuscript

Exercise in NAFLD: Just do it

Shelley E. Keating, Leon A. Adams

PII: S0168-8278(16)30317-8

DOI: <http://dx.doi.org/10.1016/j.jhep.2016.06.022>


Reference: JHEPAT 6169

To appear in: *Journal of Hepatology*

Received Date: 21 June 2016

Accepted Date: 27 June 2016

Please cite this article as: Keating, S.E., Adams, L.A., Exercise in NAFLD: Just do it, *Journal of Hepatology* (2016), doi: <http://dx.doi.org/10.1016/j.jhep.2016.06.022>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Exercise in NAFLD: Just do it.

Shelley E. Keating<sup>1</sup> & Leon A. Adams<sup>2</sup>

1. Centre for Research on Exercise, Physical Activity and Health (CRExPAH), School of Human Movement and Nutrition Sciences, The University of Queensland, Brisbane, Australia.
2. School of Medicine and Pharmacology, The University of Western Australia, Nedlands, Australia.

## Contact Details:

Shelley E. Keating, PhD  
Centre for Research on Exercise, Physical Activity and Health (CRExPAH)  
School of Human Movement and Nutrition Sciences, UQ  
HMS Building (26B)  
St Lucia, QLD 4072  
Australia

**T:** +61 7 3346 9999  
**F:** +61 7 3365 6877  
Email: s.keating@uq.edu.au

Leon A. Adams MBBS, PhD, FRACP  
M503  
School of Medicine and Pharmacology, UWA  
QEI Medical Campus  
Verdun St  
Nedlands, WA 6009  
Australia

**T:** +61 8 6151 1052  
**F:** +61 8 6151 1028  
Email: leon.adams@uwa.edu.au

Word Count: 1492    Figures: 0    Tables: 0

## Abbreviations:

**Keywords:** NAFLD, exercise, aerobic, resistance, hepatic lipid

**Conflict of Interest:** Nil

**Financial Support:** Nil

**Authors Contributions:** SEK and LAA both contributed equally to the data review and interpretation for this editorial, the manuscript writing and editing.

43 The health burden of non-alcoholic fatty liver disease (NAFLD) is multifaceted. In addition to  
44 sequelae of end-stage liver disease and hepatocellular carcinoma, the deleterious role of  
45 hepatic steatosis in the development of type 2 diabetes and cardiovascular disease is also  
46 well established.(1,2) NAFLD affects up to one third of the adult population and may  
47 become more prevalent with the escalation of obesity globally. Therefore, there is an urgent  
48 need for effective therapies for not only reducing steatosis and liver injury, but for  
49 improving insulin resistance and the cardiovascular risk associated with NAFLD.

50 Although there are many promising drug therapies in phase II and III clinical trials, treating a  
51 large proportion of the population with medications for many years will be an expensive  
52 management option at a population level. As there are no pharmacological agents currently  
53 accepted for the long-term treatment of NAFLD, the mainstay of treatment is weight loss via  
54 lifestyle intervention incorporating dietary and structured exercise intervention.(3) Indeed,  
55 weight gain is arguably the strongest determinant of the development of NAFLD (4), and  
56 weight loss is the greatest predictor of reductions in liver fat and aminotransferases.(3) Even  
57 small reductions (3-5% body weight loss) reduce steatosis and associated metabolic  
58 parameters however, larger reductions ( $\geq 10\%$ ) appear to be required for histologic  
59 improvements of liver injury.(5) Whilst weight loss is effective, it is notoriously difficult to  
60 sustain in the long term.(6) Exercise is routinely recommended for the management of  
61 NAFLD, since the efficacy of exercise *per se* for the reduction of liver fat is now recognized  
62 (7), and exercise has a multiplicity of benefits beyond weight loss. While questions regarding  
63 the optimal 'dose' of exercise therapy for reducing liver fat have been the focus of recent  
64 clinical investigations, consensus as to how effective exercise is over a longer term at a  
65 population level is unclear. Furthermore, how much exercise is required to prevent or

reverse NAFLD has remained elusive. This is in part due to the paucity of epidemiological data to support the findings of smaller clinical trials.

In this issue of the *Journal of Hepatology*, Sung and colleagues examine the association between exercise with both incident NAFLD and the resolution of NAFLD in a large cohort of adults followed over a mean of five years (8). This paper provides the first longitudinal epidemiological data to support the utility of exercise in both the prevention and treatment of NAFLD. Within the setting of an occupational health screening program, 169,347 men and women had repeat measures of liver fat (quantified with ultrasound) and physical activity. Of the 126,811 adults at baseline without NAFLD, 23% developed NAFLD at follow up and displayed an adverse cardiovascular risk profile when compared with participants who remained free from NAFLD. Of the 42,536 individuals with NAFLD at baseline, 34% of cases resolved. After adjusting for potential confounders, including change in body mass index (BMI), any level of moderate-vigorous exercise was associated with both a reduced risk of incident NAFLD and with the resolution of NAFLD. The greatest benefits were observed with exercise frequency  $\geq 5$  days per week, with a 16% reduction in NAFLD incidence and 40% increase in NAFLD resolution over the follow-up. Higher levels of exercise at baseline and increasing the frequency of weekly exercise bouts over time were also associated with a lower risk of incident NAFLD and the resolution of NAFLD.

These data have evident clinical ramifications with the authors concluding that weekly exercise of moderate-vigorous intensity independently reduces the risk of developing NAFLD, and improves the resolution of existing NAFLD. Noteworthy limitations were the diagnosis of NAFLD via ultrasound, which has limited ability to detect smaller changes in liver steatosis, the lack of dietary data and the assessment of exercise by the Korean version

89 of the International Physical Activity Questionnaire Short Form. This self-report method,  
90 which recalls physical activity behavior over a seven-day period, is prone to self-report bias  
91 and typically leads to an overestimation of physical activity levels. Nevertheless, a dose-  
92 response relationship was observed between reported exercise frequency and protection  
93 from incident NAFLD as well as future NAFLD resolution, suggesting a true 'causal'  
94 relationship. However, the IPAQ questionnaire assessment is only semi-quantitative and  
95 thus the ability of the study to determine the optimal prescriptive 'dose' of exercise  
96 (encompassing the frequency, intensity, duration and type of exercise) is limited.

97 This information gap is filled in part, by smaller, short-term, randomized controlled trials  
98 (RCT's) examining the efficacy of exercise for the reduction of liver fat. Multiple studies  
99 examining aerobic exercise, typically comprising 30-60 minutes of moderate to vigorous  
100 exercise on 3-5 days per week, have consistently demonstrated benefits with mean 10-44%  
101 relative reductions in intrahepatic lipid by magnetic resonance spectroscopy.(7) Notably, a  
102 recent RCT demonstrated no difference between low volume high-intensity and high  
103 volume low-intensity aerobic exercise, in reducing hepatic steatosis, suggesting different  
104 combinations of aerobic exercise may be equally beneficial.(9) In addition to reducing liver  
105 fat, these doses of exercise have clear extra-hepatic benefits including improvements in  
106 comorbid insulin resistance, systemic inflammation, dyslipidaemia, hypertension and  
107 endothelial dysfunction.(10) Thus, the well-established protective effect of physical activity  
108 on cardiovascular morbidity and mortality in the general population is likely to be applicable  
109 to patients with NAFLD.(11) Given that cardiovascular disease remains the leading cause of  
110 death in NAFLD patients, encouraging regular exercise should be advocated independently  
111 of its effect on liver disease.

Benefits are also seen with resistance-based exercise, although the evidence for resistance-based exercise for reducing steatosis is less consistent with large heterogeneity in the dose of resistance training employed. Whether resistance training is as efficacious in reducing hepatic fat as aerobic exercise is unclear, with one small RCT demonstrating no difference between interventions whereas a larger trial over 8 months favoured aerobic exercise.(12,13) However, resistance training improves insulin sensitivity, muscle strength and function, which are important given the recently demonstrated association between sarcopenia and risk of NAFLD.(14) In accordance with clinical recommendations for resistance exercise in cardiovascular disease risk modification, the current evidence suggests that resistance training should complement, rather than replace, aerobic exercise training.(15)

As exercise has been convincingly demonstrated to reduce hepatic steatosis, improve liver enzymes and ameliorate insulin resistance, it would be anticipated that it might also improve liver inflammation and injury in patients with NAFLD. Cross-sectional data from NAFLD patients undergoing liver biopsy, suggests that individuals who reported engaging in vigorous-intensity exercise have a lower BMI-adjusted odds for non-alcoholic steatohepatitis (NASH) and advanced fibrosis.(16) However, there are no clinical trials examining optimal doses of exercise that reverse liver injury and fibrosis. Moreover, while lifestyle intervention resulting in >7-10% weight loss has been associated with a reduction in liver injury and fibrosis (6), there is currently no evidence to suggest that exercise in isolation without concomitant weight loss can reduce NASH or fibrosis. Thus there is a need to determine the interaction between exercise and NASH, fibrosis and the associated metabolic outcomes, as well as the added benefits of exercise atop of weight loss for liver-related outcomes. A major limitation for research in this field is the invasive nature of liver

biopsy, which is not palatable for the majority of patients. Thus, future population studies incorporating non-invasive assessment of fibrosis such as transient elastography or non-invasive markers are eagerly awaited.

The large sample size included in Sung and colleagues work allows for a well-powered analysis of free-living individuals at a population level, and thus considerably strengthens the consensus that exercise is effective as a prevention and treatment strategy for NAFLD. This study also reinforces that NAFLD is a dynamic condition, with one-third of the 42,536 individuals with fatty liver at baseline having resolution after five years. Resolution was associated with a modest mean reduction in BMI ( $-0.5 \text{ kg/m}^2$ ) and increase in the number of weekly exercise sessions. Thus, relatively modest weight reduction in conjunction with increased exercise frequency has the potential to significantly impact on the prevalence of NAFLD at a population level. While most successful lifestyle interventions will be multidisciplinary (including dietitians, exercise specialists and psychologists), these findings highlight that emphasis should be placed on exercise adoption and maintenance in the primary management of NAFLD. Patients with NAFLD have been shown to understand the 'benefits of exercise' however don't participate due to low confidence and a fear of falling, highlighting the need for strategies to overcome these barriers.(17) Promising therapies include high intensity interval training (HIIT) with a recent study employing a modified HIIT protocol (combined aerobic and resistance exercise) demonstrating significant reductions in steatosis and improved cardiac function.(18) Pilot studies examining acceleration training (resistance exercise performed on a vibration platform) and hybrid training (involving the voluntary and electrical contraction of muscles) demonstrate the potential for innovative strategies to also improve liver fat in NAFLD.(19,20)

While further research will enable clinicians to titrate the dose required for individual benefit across the spectrum of NAFLD, there is now clear consensus that exercise is a 'polypill' for the management of NAFLD; we just need to learn how to get patients to take it regularly and seriously.

# References:

1. Targher G, Day CP, Bonora E. Risk of Cardiovascular Disease in Patients with Nonalcoholic Fatty Liver Disease. *New England Journal of Medicine* 2010;363:1341-1350.
2. Yki-Jarvinen H. Non-alcoholic fatty liver disease as a cause and a consequence of metabolic syndrome. *Lancet Diabetes Endocrinol* 2014;2:901-910.
3. Thoma C, Day CP, Trenell MI. Lifestyle interventions for the treatment of non-alcoholic fatty liver disease in adults: A systematic review. *J Hepatol* 2012;56:255-266.
4. Suzuki A, Lindor K, Saver JS, Lymp J, Mendes F, Muto A, Okada T, et al. Effect of changes on body weight and lifestyle in nonalcoholic fatty liver disease. *J Hepatol* 2005;43:1060-1066.
5. Vilar-Gomez E, Martinez-Perez Y, Calzadilla-Bertot L, Torres-Gonzalez A, Gra-Oramas B, Gonzalez-Fabian L, Friedman SL, et al. Weight Loss Through Lifestyle Modification Significantly Reduces Features of Nonalcoholic Steatohepatitis. *Gastroenterology* 2015;149:367-378.e365.
6. Musso G, Gambino R, Cassader M, Pagano G. A meta-analysis of randomized trials for the treatment of nonalcoholic fatty liver disease. *Hepatology* 2011;52:79-104.
7. Keating S, Hackett D, George J, Johnson N. Exercise and non-alcoholic fatty liver disease: a systematic review and meta-analysis. *J Hepatol* 2012;57:157 - 166.
8. Sung K-C, Ryu S, Lee J-Y, Kim J-Y, Wild SH, Byrne CD. Development of new fatty liver, or resolution of existing fatty liver, over 5 years of follow up: effect of exercise. *Journal of Hepatology*.
9. Keating SE, Hackett DA, Parker HM, O'Connor HT, Geroft JA, Sainsbury A, Baker MK, et al. Effect of aerobic exercise training dose on liver fat and visceral adiposity. *J Hepatol* 2015; 63: 174-82
10. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee I-M, Nieman DC, et al. Quantity and Quality of Exercise for Developing and Maintaining Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Apparently Healthy Adults: Guidance for Prescribing Exercise. *Med Sci Sports Exerc* 2011;43:1334-1359 1310.1249.
11. Berlin JA, Colditz GA. A meta-analysis of physical activity in the prevention of coronary heart disease. *Am J Epidemiol* 1990;132:612-628.
12. Bacchi E, Negri C, Targher G, Faccioli N, Lanza M, Zoppini G, Zanolini E, et al. Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 randomized trial). *Hepatology* 2013;58:1287-1295.
13. Slentz CA, Bateman LA, Willis LH, Shields AT, Tanner CJ, Piner LW, Hawk VH, et al. Effects of aerobic vs. resistance training on visceral and liver fat stores, liver enzymes, and insulin resistance by HOMA in overweight adults from STRRIDE AT/RT. *Am J Physiol Endocrinol Metab* 2011;301:E1033-1039.
14. Hong HC, Hwang SY, Choi HY, Yoo HJ, Seo JA, Kim SG, Kim NH, et al. Relationship between sarcopenia and nonalcoholic fatty liver disease: the Korean Sarcopenic Obesity Study. *Hepatology* 2014;59:1772-1778.
15. Williams MA, Haskell WL, Ades PA, Amsterdam EA, Bittner V, Franklin BA, Gulanick M, et al. Resistance Exercise in Individuals With and Without Cardiovascular Disease: 2007 Update: A Scientific Statement From the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2007;116:572-584.



16. Kistler KD, Brunt EM, Clark JM, Diehl AM, Sallis JF, Schwimmer JB, Group NCR, et al. Physical Activity Recommendations, Exercise Intensity, and Histological Severity of Nonalcoholic Fatty Liver Disease. *The American Journal of Gastroenterology* 2011;106:460-468.
17. Frith J, Day CP, Robinson L, Elliott C, Jones DE, Newton JL. Potential strategies to improve uptake of exercise interventions in non-alcoholic fatty liver disease. *J Hepatol* 2010;52:112-116.
18. Hallsworth K, Thoma C, Hollingsworth KG, Cassidy S, Anstee QM, Day CP, Trenell MI. Modified high-intensity interval training reduces liver fat and improves cardiac function in non-alcoholic fatty liver disease: a randomized controlled trial. *Clin Sci (Lond)* 2015;129:1097-1105.
19. Oh S, Shida T, Sawai A, Maruyama T, Eguchi K, Isobe T, Okamoto Y, et al. Acceleration training for managing nonalcoholic fatty liver disease: a pilot study. *Ther Clin Risk Manag* 2014;10:925-936.
20. Oh S, Maruyama T, Eguchi K, Shida T, Arai E, Isobe T, Okamoto Y, et al. Therapeutic effect of hybrid training of voluntary and electrical muscle contractions in middle-aged obese women with nonalcoholic fatty liver disease: a pilot trial. *Ther Clin Risk Manag* 2015;11:371-380.